INDEX TO AUTHORS

ÅSTRÖM, KARL E., WEBSTER, HENRY DE F., and ARNASON, BARRY G. The initial lesion in experimental allergic neuritis. A phase and electron microscopic study, 469

ABEL, CARLOS A. See GREY, ABEL, YOUNT, and KUNKEL, 1223

AGNEW, HEWES D. The effect of heterologous anti-lymphocytic serum on the small lymphocyte population of rats, 111

AISENBERG, ALAN C., and DAVIS, CAROLINE. The thymus and recovery from cyclophosphamide-induced tolerance to sheep erythrocytes, 35

— and —. The thymus and recovery of the sheep erythrocyte response in irradiated mice, 1327

ALBERTINI, RICHARD J., and BACH, FRITZ H. Quantitative assay of antigenic disparity at *HL-A*—the major histocompatibility locus in man, 639

AMOS, D. BERNARD, and BACH, FRITZ. Phenotypic expressions of the major histocompatiblity locus in man (*HL-A*). Leukocyte antigens and mixed leukocyte culture reactivity, 623

ANDRÉ-SCHWARTZ, JANINE. See LEWIS, ARMSTRONG, ANDRE-SCHWARTZ, MUFTUOGLU, BELDOTTI, and SCHWARTZ, 653

AOKI, TADAO. See HÄMMERLING, AOKI, DE HARVEN, BOYSE, and OLD, 1461

ARGYRIS, BERTIE F. Role of macrophages in immunological maturation, 459

ARMSTRONG, MARTINE Y. K. See LEWIS, ARMSTRONG, ANDRE-SCHWARTZ, MUFTUOGLU, BELDOTTI, and SCHWARTZ, 653

ARNASON, BARRY G. See ÅSTRÖM, WEBSTER, and ARNASON, 469

AUSTRIAN, ROBERT. See BORNSTEIN, SCHIFFMAN, BERNHEIMER, and AUSTRIAN, 1385

AXLINE, S. G. Isozymes of acid phosphatase in normal and Calmette-Guérin bacillus–induced rabbit alveolar macrophages, 1031

AYOUB, ELIA M. See DUDDING and AYOUB, 1081

BACH, FRITZ H. See ALBERTINI and BACH, 639

—. See AMOS and BACH, 623

BAGLIONI, C. See CIOLI and BAGLIONI, 517

BARKER, CLYDE F., and BILLINGHAM, R. E. The role of afferent lymphatics in the rejection of skin homografts, 197

BAUMINGER, SARA. See GERSHON, BAUMINGER, SELA, and FELDMAN, 223

BELDOTTI, LORRAINE. See LEWIS, ARMSTRONG, ANDRE-SCHWARTZ, MUFTUOGLU, BELDOTTI, and SCHWARTZ, 653

BERGQUIST, R. See BLOTH and BERGQUIST, 1129

BERNHEIMER, HARRIET P. See BORNSTEIN, SCHIFFMAN, BERNHEIMER, and AUSTRIAN, 1385

BILLINGHAM, R. E. See BARKER and BILLINGHAM, 197

BLOTH, B., and BERGQUIST, R. The ultrastructure of human thyroglobulin, 1129

BLYTH, JANET L. See WILSON, BLYTH, and NOWELL, 1157

BORNSTEIN, DONALD L., SCHIFFMAN, GERALD, BERNHEIMER, HARRIET P., and AUSTRIAN, ROBERT. Capsula-

tion of pneumococcus with soluble C-like (C_s) polysaccharide. I. Biological and genetic properties of C_s-pneumococcal strains, 1385
BORSOS, TIBOR. See COLTEN, GORDON, BORSOS, and RAPP, 595
BOSMAN, CESARE, and FELDMAN, JOSEPH D. Cytology of immunologic memory. A morphologic study of lymphoid cells during the anamnestic response, 293
BOYSE, EDWARD A., STOCKERT, ELISABETH, and OLD, LLOYD J. Isoantigens of the *Histocompatibility-2* and *Thymus leukemia antigen loci* of the mouse, 85
—. See HÄMMERLING, AOKI, DE HARVEN, BOYSE, and OLD, 1461
BRAUN, DIETMAR G., and KRAUSE, RICHARD M. The individual antigenic specificity of antibodies to streptococcal carbohydrates, 969
—. See KARAKAWA, BRAUN, LACKLAND, and KRAUSE, 325
BRUNO, PAULINE N. See WILEY and BRUNO, 959
BYERS, VERA S., and SERCARZ, ELI E. The X-Y-Z scheme of immunocyte maturation. V. Paralysis of memory cells, 715

CAMITTA, BRUCE M. See MEDEARIS, CAMITTA, and HEATH, 399
CIOLI, D., and BAGLIONI, C. Catabolic origin of a Bence Jones protein fragment, 517
CLARK, SAM L. Incorporation of sulfate by the mouse thymus. Its relation to secretion by medullary epithelial cells and to thymic lymphopoiesis, 927
CLIFFORD, P. See KLEIN, PEARSON, NADKARNI, NADKARNI, KLEIN, HENLE, HENLE, and CLIFFORD, 1011
CLINE, MARTIN J., and SWETT, VIRGINIA C. The interaction of human monocytes and lymphocytes, 1309
COCHRANE, C. G. See RANADIVE and COCHRANE, 605
COHN, ZANVIL A. See VAN FURTH and COHN, 415
COLTEN, HARVEY R., GORDON, JAMES M., BORSOS, TIBOR, and RAPP, HERBERT J. Synthesis of the first component of human complement in vitro, 595
CONNELL, MARY ST. JAMES. See SHEARER, CUDKOWICZ, CONNELL, and PRIORE, 437
CUDKOWICZ, G. See SHEARER, CUDKOWICZ, CONNELL, and PRIORE, 437
CUNNINGHAM, A. See NOSSAL, CUNNINGHAM, MITCHELL, and MILLER, 839

DAVID, JOHN R., and SCHLOSSMAN, STUART F. Immunochemical studies on the specificity of cellular hypersensitivity. The in vitro inhibition of peritoneal exudate cell migration by chemically defined antigens, 1451
DAVIE, JOSEPH M., and OSTERLAND, C. KIRK. Chemical characterization of glycopeptides from human γM-globulins, 699
DAVIS, CAROLINE. See AISENBERG and DAVIS, 35, 1327
DE HARVEN, ETIENNE. See HÄMMERLING, AOKI, DE HARVEN, BOYSE, and OLD, 1461
DENT, PETER B. See OLSON, DENT, RAWLS, SOUTH, MONTGOMERY, MELNICK, and GOOD, 47
DIEHL, V. See KLEIN, PEARSON, HENLE, HENLE, DIEHL, and NIEDERMAN, 1021
DONALDSON, ROBERT M. See KOPP, TRIER, MACKENZIE, AND DONALDSON, 357
—. See MACKENZIE, DONALDSON, KOPP, and TRIER, 375
DUBOS, RENÉ. See LEE and DUBOS, 753
—. See SAVAGE and DUBOS, 97
DUDDING, BURTON A., and AYOUB, ELIA M. Persistence of streptococ-

cal Group A antibody in patients with rheumatic valvular disease, 1081
DUTTON, RICHARD W. See RAIDT, MISHELL, and DUTTON, 681

EIDINGER, DAVID, KHAN, SALAHUDDIN A., and MILLAR, KENNETH G. The effect of antigenic competition on various manifestations of humoral antibody formation and cellular immunity, 1183
EVANS, GEOFFREY, PACKHAM, MARIAN A., NISHIZAWA, EDWARD E., MUSTARD, JAMES F., and MURPHY, EDMUND A. The effect of acetylsalicylic acid on platelet function, 877

FEIGEN, GEORGE A. See REITZ, PRAGER, and FEIGEN, 1401
FELDMAN, JOSEPH D. See BOSMAN and FELDMAN, 293
FELDMAN, MICHAEL. See GERSHON, BAUMINGER, SELA, and FELDMAN, 223
FREEDMAN, SAMUEL O. See KRUPEY, GOLD, and FREEDMAN, 387

GERSHON, HARRIET, BAUMINGER, SARA, SELA, MICHAEL, and FELDMAN, MICHAEL. Studies on the competence of single cells to produce antibodies of two specificities, 223
GEWURZ, H., SHIN, H. S., and MERGENHAGEN, S. E. Interactions of the complement system with endotoxic lipopolysaccharide. Consumption of each of the six terminal complement components, 1049
—. See SNYDERMAN, GEWURZ, and MERGENHAGEN, 259
GOLD, PHIL. See KRUPEY, GOLD, and FREEDMAN, 387
GOOD, ROBERT A. See OLSON, DENT, RAWLS, SOUTH, MONTGOMERY, MELNICK, and GOOD, 47
GORDON, JAMES M. See COLTEN, GORDON, BORSOS, and RAPP, 595
GREEN, IRA. Distribution of antibody-forming cells of different specificities in the lymph nodes and spleens of guinea pigs, 729
GREY, HOWARD M., ABEL, CARLOS A., YOUNT, WILLIAM J., and KUNKEL, HENRY G. A subclass of human γA-globulins (γA2) which lacks the disulfide bonds linking heavy and light chains, 1223
GUSTAFSSON, B. E. See LAGERCRANTZ, HAMMARSTRÖM, PERLMANN, and GUSTAFSSON, 1339

HÄMMERLING, ULRICH, AOKI, TADAO, DE HARVEN, ETIENNE, BOYSE, EDWARD A., and OLD, LLOYD J. Use of hybrid antibody with anti-γG and anti-ferritin specificities in locating cell surface antigens by electron microscopy, 1461
HAMMARSTRÖM, S. See LAGERCRANTZ, HAMMARSTROM, PERLMANN, and GUSTAFSSON, 1339
HEATH, EDWARD C. See MEDEARIS, CAMITTA, and HEATH, 399
HEIDELBERGER, MICHAEL, and SLODKI, MOREY E. Predicted and unpredicted cross-reactions of an acetylphosphogalactan of *Sporobolomyces* yeast, 189
HENDERSON, JAMES STUART. Endotoxin as adjuvator to the transplantation of a mouse mammary tumor, 1363
HENLE, G. See KLEIN, PEARSON, HENLE, HENLE, DIEHL, and NIEDERMAN, 1021
—. See KLEIN, PEARSON, NADKARNI, NADKARNI, KLEIN, HENLE, HENLE, and CLIFFORD, 1011
HENLE W. See KLEIN, PEARSON, HENLE, HENLE, DIEHL, and NIEDERMAN, 1021
—. See KLEIN, PEARSON, NADKARNI, NADKARNI, KLEIN, HENLE, HENLE and CLIFFORD, 1011
HENRY, CLAUDIA, and JERNE, NIELS K. Competition of 19S and 7S antigen receptors in the regulation of

the primary immune response, 133

HIRSCH, MARTIN S., NAHMIAS, ANDRÉ J., MURPHY, FREDERICK A., and KRAMER, JOEL H. Cellular immunity in vaccinia infection of mice. Anti-thymocyte serum effects on primary and secondary responsiveness, 121

HOCHSTRASSER, HARRY. See SKEGGS, LENTZ, KAHN, and HOCHSTRASSER, 13

HURD, ERIC R., and ZIFF, MORRIS. Studies on the anti-inflammatory action of 6-mercaptopurine, 785

JANOFF, A., and SCHERER, J. Mediators of inflammation in leukocyte lysosomes. IX. Elastinolytic activity in granules of human polymorphonuclear leukocytes, 1137

JERNE, NIELS K. See HENRY and JERNE, 133

KAHN, JOSEPH R. See SKEGGS, LENTZ, KAHN, and HOCHSTRASSER, 13

KARAKAWA WALTER W., BRAUN, DIETMAR G., LACKLAND, HENRY, and KRAUSE, RICHARD M. Immunochemical studies on the cross-reactivity between streptococcal and staphylococcal mucopeptide, 325

KHAN, SALAHUDDIN A. See EIDINGER, KHAN, and MILLAR, 1183

KLEIN, E. See KLEIN, PEARSON, NADKARNI, NADKARNI, KLEIN, HENLE, HENLE, and CLIFFORD, 1011

KLEIN, G., PEARSON, G., NADKARNI, J. S., NADKARNI, J. J., KLEIN, E., HENLE, G., HENLE, W., and CLIFFORD, P. Relation between Epstein-Barr viral and cell membrane immunofluorescence in Burkitt tumor cells. I. Dependence of cell membrane immunofluorescence on presence of EB virus, 1011

—, PEARSON, G., HENLE, G., HENLE, W., and NIEDERMAN, J. II. Comparison of cells and sera from patients with Burkitt's lymphoma and infectious mononucleosis, 1021

KÖLSCH, E., and MITCHISON, N. A. The subcellular distribution of antigen in macrophages, 1059

KOPP, WILLIAM L., TRIER, JERRY S., MACKENZIE, IAIN L., and DONALDSON, ROBERT M. Antibodies to intestinal microvillous membranes. I. Characterization and morphologic localization, 357

—. See MACKENZIE, DONALDSON, KOPP, AND TRIER, 375

KOROS, AURELIA M. C., MAZUR, JOHN M., and MOWERY, MARGARET J. Radioautographic studies of plaque-forming cells. I. Antigen-stimulated proliferation of plaque-forming cells, 235

KRAMER, JOEL H. See HIRSCH, NAHMIAS, MURPHY, and KRAMER, 121

KRAUSE, RICHARD M. See BRAUN and KRAUSE, 969

—. See KARAKAWA, BRAUN, LACKLAND, and KRAUSE, 325

KRUPEY, JOHN, GOLD, PHIL, and FREEDMAN, SAMUEL O. Physicochemical studies of the carcinoembryonic antigens of the human digestive system, 387

KUNKEL, HENRY G. See GREY, ABEL, YOUNT, and KUNKEL, 1223

—. See NATVIG, KUNKEL, YOUNT, and NIELSEN, 763

LACKLAND, HENRY. See KARAKAWA, BRAUN, LACKLAND, and KRAUSE, 325

LAGERCRANTZ, R., HAMMARSTRÖM, S., PERLMANN, P., and GUSTAFSSON, B. E. Immunological studies in ulcerative colitis. IV. Origin of autoantibodies, 1339

LANCE, EUGENE M. See TAUB and LANCE, 1281

LAY, WALTRAUT H., and NUSSENZ-

WEIG, VICTOR. Receptors for complement on leukocytes, 991
LEE, CHI-JEN, and DUBOS, RENÉ. Lasting biological effects of early environmental influences. III. Metabolic responses of mice to neonatal infection with a filterable weight-depressing agent, 753
LENTZ, KENNETH E. See SKEGGS, LENTZ, KAHN, and HOCHSTRASSER, 13
LEVY, L. See VAN WINKLE and LEVY, 497
LEWIS, ROBERT M., ARMSTRONG, MARTINE Y. K., ANDRÉ-SCHWARTZ, JANINE, MUFTUOGLU, ASUMAN, BELDOTTI, LORRAINE, and SCHWARTZ, ROBERT S. Chronic allogeneic disease. I. Development of glomerulonephritis, 653
LUBAROFF, DAVID M., and WAKSMAN, BYRON H. Bone marrow as source of cells in reactions of cellular hypersensitivity. I. Passive transfer of tuberculin sensitivity in syngeneic systems, 1425
— and —. II. Identification of allogeneic or hybrid cells by immunofluorescence in passively transferred tuberculin reactions, 1427

MACKENZIE, IAIN L., DONALDSON, ROBERT M., KOPP, WILLIAM L., and TRIER, JERRY S. Antibodies to intestinal microvillous membranes. II. Inhibition of intrinsic factor–mediated attachment of vitamin B_{12} to hamster brush borders, 375
—. See KOPP, TRIER, MACKENZIE, and DONALDSON, 357
MALAVIYA, ANAND N. See TANNENBERG and MALAVIYA, 895
MARTIN, W. J., and MILLER, J. F. A. P. Cell to cell interaction in the immune response. IV. Site of action of antilymphocyte globulin, 855
MAZUR, JOHN M. See KOROS, MAZUR, and MOWERY, 235
MCDEVITT, HUGH O., and TYAN, MARVIN L. Genetic control of the antibody response in inbred mice. Transfer of response by spleen cells and linkage to the major histocompatibility (*H-2*) locus, 1
MEDEARIS, DONALD N., CAMITTA, BRUCE M., and HEATH, EDWARD C. Cell wall composition and virulence in *Escherichia coli*, 399
MELNICK, JOSEPH L. See OLSON, DENT, RAWLS, SOUTH, MONTGOMERY, MELNICK, and GOOD, 47
MERGENHAGEN, S. E. See GEWURZ, SHIN, and MERGENHAGEN, 1049
—. See SNYDERMAN, GEWURZ, and MERGENHAGEN, 259
MESSNER, RONALD P. See QUIE, MESSNER, and WILLIAMS, 553
MEYER, KARL. See SHULMAN and MEYER, 1353
MILLAR, KENNETH G. See EIDINGER, KHAN, and MILLAR, 1183
MILLER, J. F. A. P., and MITCHELL, G. F. Cell to cell interaction in the immune response. I. Hemolysin-forming cells in neonatally thymectomized mice reconstituted with thymus or thoracic duct lymphocytes, 801
—. See MARTIN and MILLER, 855
—. See MITCHELL and MILLER, 821
—. See NOSSAL, CUNNINGHAM, MITCHELL, and MILLER, 839
MISHELL, R. I. See RAIDT, MISHELL, and DUTTON, 681
MITCHELL, G. F., and MILLER, J. F. A. P. Cell to cell interaction in the immune response. II. The source of hemolysin-forming cells in irradiated mice given bone marrow and thymus or thoracic duct lymphocytes, 821
—. See MILLER and MITCHELL, 801
—. See NOSSAL, CUNNINGHAM, MITCHELL, and MILLER, 839

MITCHISON, N. A. See KÖLSCH and MITCHISON, 1059
MONTGOMERY, J. R. See OLSON, DENT, RAWLS, SOUTH, MONTGOMERY, MELNICK, and GOOD, 47
MOWERY, MARGARET J. See KOROS, MAZUR, and MOWERY, 235
MÜLLER-EBERHARD, HANS J. See POLLEY and MÜLLER-EBERHARD, 533
MUFTUOGLU, ASUMAN. See LEWIS, ARMSTRONG, ANDRE-SCHWARTZ, MUFTUOGLU, BELDOTTI, and SCHWARTZ, 653
MURPHY, EDMUND A. See EVANS, PACKHAM, NISHIZAWA, MUSTARD, and MURPHY, 877
MURPHY, FREDERICK A. See HIRSCH, NAHMIAS, MURPHY, and KRAMER, 121
MUSTARD, JAMES F. See EVANS, PACKHAM, NISHIZAWA, MUSTARD, and MURPHY, 877

NADKARNI, J. J. See KLEIN, PEARSON, NADKARNI, NADKARNI, KLEIN, HENLE, HENLE, and CLIFFORD, 1011
NADKARNI, J. S. See KLEIN, PEARSON, NADKARNI, NADKARNI, KLEIN, HENLE, HENLE, and CLIFFORD, 1011
NAFF, GEORGE B., and RATNOFF, OSCAR D. The enzymatic nature of C′1r. Conversion of C′1s to C′1 esterase and digestion of amino acid esters by C′1r, 571
NAHMIAS, ANDRÉ J. See HIRSCH, NAHMIAS, MURPHY, and KRAMER, 121
NATVIG, J. B., KUNKEL, HENRY G., YOUNT, W. J., and NIELSEN, J. C. Further studies on the γG–heavy chain gene complexes, with particular reference to the genetic markers Gm(g) and Gm(n), 763
NIEDERMAN, J. C. See KLEIN, PEARSON, HENLE, HENLE, DIEHL, and NIEDERMAN, 1021
NIELSEN, J. C. See NATVIG, KUNKEL, YOUNT, and NIELSEN, 763
NISHIZAWA, EDWARD E. See EVANS, PACKHAM, NISHIZAWA, MUSTARD, and MURPHY, 877
NOSSAL, G. J. V., CUNNINGHAM, A., MITCHELL, G. F., and MILLER, J. F. A. P. Cell to cell interaction in the immune response. III. Chromosomal marker analysis of single antibody-forming cells in reconstituted, irradiated, or thymectomized mice, 839
NOWELL, PETER C. See WILSON, BLYTH, and NOWELL, 1157
NUSSENZWEIG, VICTOR. See LAY and NUSSENZWEIG, 991

OLD, LLOYD J. See BOYSE, STOCKERT, and OLD, 85
—. See HÄMMERLING, AOKI, DE HARVEN, BOYSE, and OLD, 1461
OLSON, GEORGE B., DENT, PETER B., RAWLS, WILLIAM E., SOUTH, MARY ANN, MONTGOMERY, J. R., MELNICK, JOSEPH L., and GOOD, ROBERT A. Abnormalities of in vitro lymphocyte responses during rubella virus infections, 47
OSTERLAND, C. KIRK. See DAVIE and OSTERLAND, 699
OWENS, ALBERT H., and SANTOS, GEORGE W. The induction of graft versus host disease in mice treated with cyclophosphamide, 277

PACKHAM, MARIAN A. See EVANS, PACKHAM, NISHIZAWA, MUSTARD, and MURPHY, 877
PEARSON, G. See KLEIN, PEARSON, HENLE, HENLE, DIEHL, and NIEDERMAN, 1021
—. See KLEIN, PEARSON, NADKARNI, NADKARNI, KLEIN, HENLE, HENLE, and CLIFFORD, 1011
PERLMANN, P. See. LAGERCRANTZ, HAMMARSTRÖM, PERLMANN, and GUSTAFSSON, 1339
POLLEY, MARGARET J., and MÜLLER-EBERHARD, HANS J. The second

component of human complement. Its isolation, fragmentation by C'1 esterase, and incorporation into C'3 convertase, 533
PRAGER, DENIS J. See REITZ, PRAGER, and FEIGEN, 1401
PRIORE, R. L. See SHEARER, CUDKOWICZ, CONNELL, and PRIORE, 437

QUIE, PAUL G., MESSNER, RONALD P., and WILLIAMS, RALPH C., JR. Phagocytosis in subacute bacterial endocarditis. Localization of the primary opsonic site to Fc fragment, 553

RAIDT, DONALD J., MISHELL, R. I., and DUTTON, RICHARD W. Cellular events in the immune response. Analysis and in vitro response of mouse spleen cell populations separated by differential flotation in albumin gradients, 681
RANADIVE, N. S., and COCHRANE, C. G. Isolation and characterization of permeability factors from rabbit neutrophils, 605
RAPP, HERBERT J. See COLTEN, GORDON, BORSOS, and RAPP, 595
RATNOFF, OSCAR D. See NAFF and RATNOFF, 571
RAWLS, WILLIAM E. See OLSON, DENT, RAWLS, SOUTH, MONTGOMERY, MELNICK, and GOOD, 47
REITZ, BRUCE A., PRAGER, DENIS J., and FEIGEN, GEORGE A. An analysis of the toxic actions of purified streptolysin O on the isolated heart and separate cardiac tissues of the guinea pig, 1401
RICHTER, MAXWELL. See SINGHAL and RICHTER, 1099
ROOT RICHARD K., and WOLFF, SHELDON M. Pathogenic mechanisms in experimental immune fever, 309
RUDDLE, NANCY H., and WAKSMAN, BYRON H. Cytotoxicity mediated by soluble antigen and lymphocytes in delayed hypersensitivity. I. Characterization of the phenomenon, 1237
— and —. II. Correlation of the in vitro response with skin reactivity, 1255
— and —. III. Analysis of mechanism, 1267

SANTOS, GEORGE W. See OWENS and SANTOS, 277
SAVAGE, DWAYNE C., and DUBOS, RENÉ. Alterations in the mouse cecum and its flora produced by antibacterial drugs, 97
SCHERER, J. See JANOFF and SCHERER, 1137
SCHIFFMAN, GERALD. See BORNSTEIN, SCHIFFMAN, BERNHEIMER, and AUSTRIAN, 1385
SCHLOSSMAN, STUART F. See DAVID and SCHLOSSMAN, 1451
SCHWARTZ, ROBERT S. See LEWIS, ARMSTRONG, ANDRÉ-SCHWARTZ, MUFTUOGLU, BELDOTTI, and SCHWARTZ, 653
SELA, MICHAEL. See GERSHON, BAUMINGER, SELA, and FELDMAN, 223
SELL, STEWART. Studies on rabbit lymphocytes in vitro. IX. The suppression of antiallotype-induced blast transformation in lymphocyte cultures from allotypically suppressed donors, 341
SERCARZ, ELI E. See BYERS and SERCARZ, 715
SHEARER, G. M., CUDKOWICZ, G. CONNELL, MARY ST. JAMES, and PRIORE, R. L. Cellular differentiation of the immune system of mice. I. Separate splenic antigen-sensitive units for different types of anti-sheep antibody-forming cells, 437
SHIN, H. S. See GEWURZ, SHIN, and MERGENHAGEN, 1049

SHULMAN, HERBERT J., and MEYER, KARL. Cellular differentiation and the aging process in cartilaginous tissues. Mucopolysaccharide synthesis in cell cultures of chondrocytes, 1353

SILVERS, WILLYS K. Studies on the induction of tolerance of the H-Y antigen in mice with neonatal skin grafts, 69

SINGHAL, SHARWAN K., and RICHTER, MAXWELL. Cells involved in the immune response. IV. The response of normal and immune bone marrow and lymphoid tissue lymphocytes to antigens in vitro, 1099

SKEGGS, LEONARD T., LENTZ, KENNETH E., KAHN, JOSEPH R., and HOCHSTRASSER, HARRY. Kinetics of the reaction of renin with nine synthetic peptide substrates, 13

SLODKI, MOREY E. See HEIDELBERGER and SLODKI, 189

SNYDERMAN, R., GEWURZ, H., and MERGENHAGEN, S. E. Interactions of the complement system with endotoxic lipopolysaccharide. Generation of a factor chemotactic for polymorphonuclear leukocytes, 259

SOUTH, MARY ANN. See OLSON, DENT, RAWLS, SOUTH, MONTGOMERY, MELNICK, and GOOD, 47

STOCKERT, ELISABETH. See BOYSE, STOCKERT, and OLD, 85

SWETT, VIRGINIA C. See CLINE and SWETT, 1309

TANNENBERG, WALTER J. K., and MALAVIYA, ANAND N. The life cycle of antibody-forming cells. I. The generation time of 19S hemolytic plaque-forming cells during the primary and secondary responses, 895

TAUB, ROBERT N., and LANCE, EUGENE M. Histopathological effects in mice of heterologous anti-lymphocyte serum, 1281

THORBECKE, G. J. See WAKEFIELD and THORBECKE, 153, 171

TRIER, JERRY S. See KOPP, TRIER, MACKENZIE, and DONALDSON, 357

—. See MACKENZIE, DONALDSON, KOPP, and TRIER, 375

TYAN, MARVIN L. See MCDEVITT and TYAN, 1

VAN FURTH, RALPH, and COHN, ZANVIL A. The origin and kinetics of mononuclear phagocytes, 415

VAN WINKLE, M., and LEVY, L. Effect of removal of cholesterol diet upon serum sickness–cholesterol-induced atherosclerosis, 497

WAKEFIELD, J. D., and THORBECKE, G. J. Relationship of germinal centers in lymphoid tissue to immunological memory. I. Evidence for the formation of small lymphocytes upon transfer of primed splenic white pulp to syngeneic mice, 153

— and —. II. The detection of primed cells and their proliferation upon cell transfer to lethally irradiated syngeneic mice, 171

WAKSMAN, BYRON H. See LUBAROFF and WAKSMAN, 1425, 1437

—. See RUDDLE and WAKSMAN, 1237, 1255, 1267

WARD, PETER A. Chemotaxis of mononuclear cells, 1201

WEBSTER, HENRY DE F. See ÅSTRÖM, WEBSTER, and ARNASON, 469

WILEY, GROVE G., and BRUNO, PAULINE N. Cross-reactions among Group A streptococci. I. Precipitin and bacterial cross-reactions among types 33, 41, 43, 52, and Ross, 959

WILLIAMS, RALPH C., JR. See QUIE, MESSNER, and WILLIAMS, 553

WILSON, DARCY B., BLYTH, JANET L., and NOWELL, PETER C. Quantitative studies on the mixed lymphocyte interaction in rats. III. Kinetics of the response, 1157
WOLFF, SHELDON M. See ROOT and WOLFF, 309
YOUNT, WILLIAM J. See GREY, ABEL, YOUNT, and KUNKEL, 1223
—. See NATVIG, KUNKEL, YOUNT, and NIELSEN, 763
ZIFF, MORRIS. See HURD and ZIFF, 785

INDEX TO SUBJECTS

ABNORMALITIES of in vitro lymphocyte responses during rubella virus infections, 47

Acetylphosphogalactan of *Sporobolomyces* yeast, cross-reactions of, 189

Acetylsalicylic acid, effect of on platelet function, 877

Acid, acetylsalicylic, effect of on platelet function, 877

—, amino, esters, digestion of by C′1r and conversion of C′1s to C′1 esterase, 571

— phosphatase, isozymes of, in normal and Calmette-Guérin bacillus-induced alveolar macrophages, 1031

Afferent lymphatics, role of, in rejection of skin homografts, 197

Aging process and cellular differentiation in cartilaginous tissues, mucopolysaccharide synthesis in chondrocyte cell cultures, 1353

Albumin gradients, analysis and in vitro response of mouse spleen cell populations separated by differential flotation in albumin gradients, 681

Allergic neuritis, initial lesion in, phase and electron microscopic study, 469

Allogeneic disease, chronic, development of glomerulonephritis, 653

— or hybrid cells, identification of by immunofluorescence in passively transferred tuberculin reactions, 1437

Allotypically suppressed donors, suppression of antiallotype-induced blast transformation in lymphocyte cultures from, 341

Alveolar macrophages, normal and Calmette-Guérin bacillus-induced, isozymes of acid phosphatase in, 1031

Amino acid esters, digestion of by C′1r and conversion of C′1s to C′1 esterase, 571

Anamnestic response, morphologic study of lymphoid cells during, 293

Antiallotype-induced blast transformation in lymphocyte cultures from allotypically suppressed donors, suppression of, 341

Antibacterial drugs, alterations produced in mouse cecum and flora by, 97

Antibodies, competence of single cells to produce antibodies of two specificities, 223

— to intestinal microvillous membranes, characterization and morphologic localization, 357

— to intestinal microvillous membranes, inhibition of intrinsic factor-mediated attachment of vitamin B_{12}, 375

— to streptoccal carbohydrates, individual antigenic specificity of, 969

Antibody formation, humoral, and cellular immunity, effect of antigenic competition on, 1183

— response, genetic control of, transfer of response by spleen cells and linkage to major histocompatibility locus, 1

—, streptococcal Group A, persistence of, in patients with rheumatic valvular disease, 1081

—, use of hybrid, with anti-γG and anti-ferritin specificities in locating

cell surface antigens by electron microscopy, 1461
Antibody-forming cells, distribution of, of different specificities, in the lymph nodes and spleen, 729
— cells, life cycle of, generation time of 19S hemolytic plaque-forming cells during primary and secondary responses, 895
— cells, separate splenic antigen-sensitive units for different types of, 437
— cells, single, chromosomal marker analysis of, 839
Antigen, H-Y, tolerance of in mice with neonatal skin grafts, 69
— receptors, competition of 19S and 7S, in regulation of primary immune response, 133
—, soluble, and lymphocytes, cytotoxicity mediated by, in delayed hypersensitivity, analysis of, 1267
—, soluble, and lymphocytes, cytotoxicity mediated by, in delayed hypersensitivity, characterization of, 1237
—, soluble, and lymphocytes, cytotoxicity mediated by, in delayed hypersensitivity, correlation of in vitro response with skin reactivity, 1255
—, subcellular distribution of, in macrophages, 1059
Antigen-sensitive units, separate splenic, for different types of anti-sheep antibody-forming cells, 437
Antigen-stimulated proliferation of plaque-forming cells, radioautographic studies, 235
Antigenic competition, effect of, on humoral antibody formation and cellular immunity, 1183
— disparity, quantitative assay of, at major histocompatibility locus in man, 639
— specificity, individual, of antibodies to streptococcal carbohydrates, 969
Antigens, carcinoembryonic, of human digestive system, physicochemical studies of, 387
—, cell surface, use of hybrid antibody with anti-γG and anti-ferritin specificities in locating by electron microscopy, 1461
—, chemically defined, in vitro inhibition of peritoneal exudate cell migration by, 1451
— in vitro, response of normal and immune bone marrow and lymphoid tissue lymphocytes to, 1099
—, leukocyte, and mixed leukocyte culture reactivity, phenotypic expressions of major histocompatibility locus in man, 623
Anti-sheep antibody-forming cells, separate splenic antigen-sensitive units for different types of, 437
Atherosclerosis, serum sickness–cholesterol-induced, effect of removal of cholesterol diet upon, 497
Autoantibodies, origin of, immunological studies of ulcerative colitis, 1339

BACILLUS-induced, Calmette-Guérin, and normal alveolar macrophages, isozymes of acid phosphate in, 1031
Bacterial endocarditis, subacute, phagocytosis in, localization of primary opsonic site to Fc fragment, 553
Bactericidal and precipitin cross-reactions among Group A streptococci, 959
Bence Jones protein fragment, catabolic origin of, 517
Biological effects, lasting, of early environmental influences, metabolic responses to neonatal infection with filterative weight-depressing agent, 753
Blast transformation, antiallotype-induced, in lymphocyte cultures from allotypically suppressed donors, suppression of, 341

Bone marrow as a source of cells in reactions of cellular hypersensitivity, identification of allogeneic or hybrid cells by immunofluorescence in passively transferred tuberculin reactions, 1437
— marrow as a source of cells in reactions of cellular hypersensitivity, passive transfer of tuberculin sensitivity in syngeneic systems, 1425
— marrow and lymphoid tissue lymphocytes, normal and immune response to antigens in vitro, 1099
— marrow, source of hemolysin-forming cells in irradiated mice given thoracic duct lymphocytes or thymus and, 821
Brush borders, hamster, inhibition of intrinsic factor–mediated attachment of vitamin B_{12} to, 375
Burkitt tumor cells, comparison with infectious mononuclear cells in Epstein-Barr viral and cell membrane immunofluorescence, 1021
— tumor cells, dependence of cell membrane immunofluorescence of on presence of Epstein-Barr virus, 1011

CALMETTE-GUERIN bacillus–induced and normal alveolar macrophages, isozymes of acid phosphatase in, 1031
Carbohydrates, streptococcal, individual antigenic specificity of antibodies to, 969
Carcinoembryonic antigens of human digestive system, physicochemical studies of, 387
Cardiac tissues, separate, and isolated heart, analysis of toxic actions of streptolysin O on, 1401
Cartilaginous tissues, cellular differentiation and aging process in, mucopolysaccharide synthesis in chondrocyte cell cultures, 1353
Catabolic origin of a Bence Jones protein fragment, 517
Cecum and flora, mouse, alterations produced in by antibacterial drugs, 97
Cell cultures, chondrocyte, mucopolysaccharide synthesis in, cellular differentiation and aging process in cartilaginous tissues, 1353
— membrane and Epstein-Barr viral immunofluorescence, comparison of cells and sera from patients with Burkitt's lymphoma and infectious mononucleosis, 1021
— membrane immunofluorescence of Burkitt tumor cells, dependence of on presence of Epstein-Barr virus, 1011
— migration, peritoneal exudate, in vitro inhibition of by chemically defined antigens, 1451
— populations, spleen, separated by differential flotation in albumin gradients, analysis and in vitro responses, 681
— surface antigens, use of hybrid antibody with anti-γG and anti-ferritin specificities in locating by electron microscopy, 1461
— to cell interaction in the immune response, chromosomal marker analysis of single antibody-forming cells, 839
— to cell interaction in immune response, hemolysin-forming cells in neonatally thymectomized mice reconstituted with thymus or thoracic duct lymphocytes, 801
— to cell interaction in immune response, site of action of anti-lymphocyte globulin, 855
— to cell interaction in immune response, source of hemolysin-forming cells in irradiated mice given bone marrow and thymus or thoracic duct lymphocytes, 821
— transfer, to lethally irradiated syngeneic mice, detection of primed cells and their proliferation upon, 171
— wall composition and virulence in *Escherichia coli*, 399

Cells, allogeneic or hybrid, identification of by immunofluorescence in passively transferred tuberculin reactions, 1437
—, antibody-forming, life cycle of, generation time of 19S hemolytic plaque-forming cells during primary and secondary responses, 895
—, antibody-forming, of different specificities, distribution of in lymph nodes and spleen, 729
—, anti-sheep antibody-forming, separate splenic antigen-sensitive units for different types of, 437
—, bone marrow as a source of, in reactions of cellular hypersensitivity, identification of allogeneic or hybrid cells by immunofluorescence in passively transferred tuberculin reactions, 1437
—, bone marrow as a source of, in reactions of cellular hypersensitivity, passive transfer of tuberculin sensitivity in syngeneic systems, 1425
—, Burkitt tumor, comparison with infectious mononuclear cells in Epstein-Barr viral and cell membrane immunofluorescence, 1021
—, Burkitt tumor, dependence of cell membrane immunofluorescence of on presence of Epstein-Barr virus, 1011
—, hemolysin-forming, in neonatally thymectomized mice reconstituted with thymus or thoracic duct lymphocytes, 801
—, hemolysin-forming, source of, in irradiated mice given bone marrow and thymus or thoracic duct lymphocytes, 821
—, 19S hemolytic plaque-forming, generation time during primary and secondary responses, 895
— involved in immune response, normal and immune bone marrow and lymphoid tissue lymphocytes and antigens in vitro, 1099
—, lymphoid, morphologic study of during anamnestic response, 293
Cells, medullary epithelial, secretion by, and thymic lymphopoiesis, relation to sulfate incorporation by thymus, 927
—, memory, paralysis of, X-Y-Z scheme of immunocyte maturation, 715
—, mononuclear, chemotaxis of, 1201
—, plaque-forming, radioautographic studies of antigen-stimulated proliferation of, 235
—, primed, detection of and proliferation of, upon transfer to lethally irradiated syngeneic mice, 171
—, single antibody-forming, chromosomal marker analysis of, 839
—, single, competence of to produce antibodies of two specificities, 223
—, spleen, transfer of antibody response by, and linkage to major histocompatibility locus, 1
Cellular differentiation and aging process in cartilaginous tissues, mucopolysaccharide synthesis in chondrocyte cell cultures, 1353
— differentiation of immune system, separate splenic antigen-sensitive units for different types of anti-sheep antibody-forming cells, 437
— events in the immune response, analysis and in vitro response of mouse spleen cell populations separated by differential flotation in albumin gradients, 681
— hypersensitivity, bone marrow as a source of cells in, identification of allogeneic or hybrid cells by immunofluorescence in passively transferred tuberculin reactions, 1437
— hypersensitivity, bone marrow as a source of cells in, passive transfer of tuberculin sensitivity in syngeneic systems, 1425
— hypersensitivity, specificity of, in

vitro inhibition of peritoneal exudate cell migration by chemically defined antigens, 1451
Cellular immunity and humoral antibody formation, effect of antigenic competition on, 1183
— immunity in vaccinia infection, 121
Chain, γG-heavy, gene complexes, with reference to genetic markers Gm(g) and Gm(n), 763
Chains, heavy and light, subclass of human γA-globulin (γA2) lacking disulfide bonds linking, 1223
Chemotactic factor, generation of, for polymorphonuclear leukocytes, 259
Chemotaxis of mononuclear cells, 1201
Cholesterol diet, effect of removal upon serum sickness–cholesterol-induced atherosclerosis, 497
Cholesterol-induced atherosclerosis, effect of removal of cholesterol diet upon serum sickness, 497
Chondrocyte cell cultures, mucopolysaccharide synthesis in, cellular differentiation and aging process in cartilaginous tissues, 1353
Chromosomal marker analysis of single antibody-forming cells, 839
Colitis, ulcerative, immunological studies of, origin of autoantibodies, 1339
Complement components, terminal, consumption of, interactions of complement system with endotoxic lipopolysaccharide, 1049
—, enzymatic nature of C′1r, conversion of C′1s to C′1 esterase and digestion of amino acid esters by C′1r, 571
—, human, second component of, its isolation, fragmentation by C′1 esterase, and incorporation into C′3 convertase, 533
Complement, human, synthesis of first component of, in vitro, 595
—, receptors for, on leukocytes, 991
— system, interactions of, with endotoxic lipopolysaccharide, consumption of terminal complement components, 1049
— system, interactions with endotoxic lipopolysaccharide, generation of factor chemotactic for polymorphonuclear leukocytes, 259
Component, first, of human complement in vitro, synthesis, of, 595
—, second, of human complement, isolation, fragmentation by C′1 esterase, and incorporation into C′3 convertase, 533
Components, terminal complement, consumption of, interactions of complement system with endotoxic lipopolysaccharide, 1049
Convertase, C′3, incorporation of second component of human complement into, 533
Cross-reactions, precipitin and bactericidal, among Group A streptococci, 959
—, predicted and unpredicted, of acetylphosphogalactan of *Sporobolomyces* yeast, 189
Cross-reactivity between streptoccal and staphylococcal mucopeptide, immunochemical studies, 325
Culture reactivity, mixed leukocyte, and leukocyte antigens, phenotypic expressions of major histocompatibility locus in man, 623
Cultures, cell, of chondrocytes, mucopolysaccharide synthesis in, 1353
—, lymphocyte, from allotypically suppressed donors, suppression of antiallotype-induced blast transformation in, 341
Cyclophosphamide, graft versus host disease in mice treated with, 277
Cyclophosphamide-induced tolerance to sheep erythrocytes, in thymus and recovery from, 35
Cytology of immunologic memory,

morphologic study of lymphoid cells during anamnestic response, 293
Cytotoxicity mediated by soluble antigen and lymphocytes in delayed hypersensitivity, analysis of, 1267
— mediated by soluble antigen and lymphocytes in delayed hypersensitivity, characterization of, 1237
— mediated by soluble antigen and lymphocytes in delayed hypersensitivity, correlation of in vitro response with skin reactivity, 1255

DIET, cholesterol, effect of removal upon serum sickness–cholesterol-induced atherosclerosis, 497
Digestion of amino acid esters by C′1r and conversion of C′1s to C′1 esterase, 571
Digestive system, human, carcinoembryonic antigens of, 387
Disease, chronic allogeneic, development of glomerulonephritis, 653
—, rheumatic valvular, persistence of streptococcal Group A antibody in patients with, 1081
Disulfide bonds linking heavy and light chains, subclass of human γA-globulin (γA2) lacking, 1223
Drugs, antibacterial, alterations produced in mouse cecum and flora by, 97

ELASTINOLYTIC activity in granules of human polymorphonuclear leukocytes, 1137
Electron and microscopic study of initial lesion in allergic neuritis, 469
— microscopy, use of hybrid antibody with anti-γG and anti-ferritin specificities in locating cell surface antigens by, 1461
Embryonic, carcino-, studies of human digestive system, 387
Endocarditis, subacute bacterial, phagocytosis in, localization of primary opsonic site to Fc fragment, 553
Endotoxic lipopolysaccharide, interactions with complement system, consumption of terminal complement components, 1049
— lipopolysaccharide, interactions with complement system, generation of chemotactic factor, 259
Endotoxin as adjuvator to transplantation of mouse mammary tumor, 1363
Environmental influences, early, lasting biological effects of, metabolic responses to neonatal infection with filterable weight-depressing agent, 753
Enzymatic nature of C′1r, conversion of C′1s to C′1 esterase and digestion of amino acid esters by C′1r, 571
Epithelial cells, medullary, secretion by, and thymic lymphopoiesis, relation to sulfate incorporation by thymus, 927
Epstein-Barr viral and cell membrane immunofluorescence, comparison of cells and sera from patients with Burkitt's lymphoma and infectious mononucleosis, 1021
— virus, dependence of cell membrane immunofluorescence of Burkitt tumor cells on presence of, 1011
Erythrocyte response, sheep, thymus and recovery of in irradiated mice, 1327
Erythrocytes, the thymus and recovery from cyclophosphamide-induced tolerance to, 35
Escherichia coli, cell wall composition and virulence in, 399
Esterase, C′1, fragmentation of second

component of human complement by, 533
Esterase, conversion of C′1s to C′1, and digestion of amino acid esters by C′1r, 571
Esters, amino acid, digestion of by C′1r and conversion of C′1s to C′1 esterase, 571

FACTORS, permeability, from neutrophils, isolation and characterization of, 605
Ferritin, anti-, and anti-γG specificities, use of hybrid antibody with in locating cell surface antigens by electron microscopy, 1461
Fever, immune, pathogenetic mechanisms in, 309
Flora and cecum, mouse, alterations produced in by antibacterial drugs, 97
Flotation, differential, in albumin gradients, analysis and in vitro response of spleen cell populations separated by, 681
Fragment, Bence Jones protein, catabolic origin of, 517
—, Fc, localization of primary opsonic site to, phagocytosis in subacute bacterial endocarditis, 553

GENE complexes, γG-heavy chain, with reference to genetic markers Gm(g) and Gm(n), 763
Genetic control of antibody response, transfer of response by spleen cells and linkage to major histocompatibility locus, 1
— markers Gm(g) and Gm(n), γG–heavy chain gene complexes with reference to, 763
Germinal centers in lymphoid tissue, relationship to immunological memory, 153, 171
Globulin, anti-lymphocyte, site of action, 855
—, thyro-, human, ultrastructure of, 1129
γA-Globulin, human, a subclass of (γA2) which lacks the disulfide bonds linking heavy and light chains, 1223
γG-Globulin, anti-, and anti-ferritin specificities, use of hybrid antibody with in locating cell surface antigens by electron microscopy, 1461
γM-Globulins, human, chemical characterization of glycopeptides from, 699
Glomerulonephritis, development of, chronic allogeneic disease, 653
Glycopepetides, chemical characterization of, from human γM-globulins, 699
Graft versus host disease in mice treated with cyclophosphamide, 277
Grafts, neonatal skin, tolerance of H-Y antigen in mice with, 69

HEART, isolated, and separate cardiac tissues, analysis of toxic action of purified streptolysin O on, 1401
γG–heavy chain gene complexes, with particular reference to genetic markers Gm(g) and Gm(n), 763
Hemolysin-forming cells in neonatally thymectomized mice reconstituted with thymus or thoracic duct lymphocytes, 801
— cells, source of, in irradiated mice given bone marrow and thymus or thoracic duct lymphocytes, 821
Hemolytic plaque-forming cells, 19S, generation time during primary and secondary responses, 895
Heterologous anti-lymphocyte serum, effect on small lymphocyte population, 111
— anti-lymphocyte serum, histopathological effects of, 1281
Histocompatibility locus, major, in man, phenotypic expression of, leukocyte antigens and mixed leukocyte culture reactivity, 623

— locus, major, in man, quantitative assay of antigenic disparity at, 639
— locus, major, transfer of antibody response by spleen cells and linkage to, 1
Histocompatibility-2 (*H-2*) and *Thymus leukemia antigen* (*Tla*) loci, isoantigens of, 85
Histopathological effects in mice of heterologous anti-lymphocyte serum, 1281
Homografts, skin, role of afferent lymphatics in rejection of, 197
Host disease versus graft in mice treated with cyclophosphamide, 277
Humoral antibody formation and cellular immunity, effect of antigenic competition on, 1183
Hypersensitivity, cellular, bone marrow as a source of cells in, identification of allogeneic or hybrid cells by immunofluorescence in passively transferred tuberculin reactions, 1437
—, cellular, bone marrow as a source of cells in, passive transfer of tuberculin sensitivity in syngeneic systems, 1425
—, cellular, specificity of, in vitro inhibition of peritoneal exudate cell migration by chemically defined antigens, 1451
—, delayed, cytotoxicity mediated by soluble antigen and lymphocytes in, analysis of, 1267
—, delayed, cytotoxicity mediated by soluble antigen and lymphocytes in, characterization of, 1237
—, delayed, cytotoxicity mediated by soluble antigen and lymphocytes in, correlation of in vitro response with skin reactivity, 1255

IMMUNE fever, pathogenetic mechanisms in, 309
— response, cell to cell interaction in, chromosomal marker analysis of single antibody-forming cells, 839
— response, cell to cell interaction in, hemolysin-forming cells in neonatally thymectomized mice reconstituted with thymus or thoracic duct lymphocytes, 801
— response, cell to cell interaction in, site of action of anti-lymphocyte globulin, 855
— response, cell to cell interaction in, source of hemolysin-forming cells in irradiated mice given bone marrow and thymus or thoracic duct lymphocytes, 821
— response, cellular events in, analysis and in vitro response of mouse spleen cell populations separated by differential flotation in albumin gradients, 681
—response, of normal and immune bone marrow and lymphoid tissue lymphocytes to antigens in vitro, 1099
— response, primary, competition of 19S and 7S antigen receptors in regulation of, 133
— system, cellular differentiation of, separate splenic antigen-sensitive units for different types of anti-sheep antibody-forming cells, 437
Immunity, cellular, and humoral antibody formation, effect of antigenic competition on, 1183
—, cellular, in vaccinia infection, 121
Immunocyte maturation, X-Y-Z scheme of, paralysis of memory cells, 715
Immunofluorescence, cell membrane, of Burkitt tumor cells, dependence of, on presence of Epstein-Barr virus, 1011
—, Epstein-Barr viral and cell membrane, comparison of cells and sera from patients with Burkitt's lymphoma and infectious mononucleosis, 1021
—, identification of allogeneic or

hybrid cells by, in passively transferred tuberculin, 1437
Immunological maturation, role of macrophages in, 459
— memory, cytology of in morphologic study of lymphoid cells during anamnestic response, 293
— memory, relationship of germinal centers in lymphoid tissue to, 153, 171
Infection, neonatal, with filterable weight-depressing agent, metabolic responses to, lasting biological effects of early environment influences, 753
—, vaccinia, cellular immunity in, 121
Infections, rubella virus, abnormalities of in vitro lymphocyte responses during, 47
Infectious mononucleosis and Burkitt's lymphoma, comparison of cells and sera from patients with, in Epstein-Barr viral and cell membrane immunofluorescence, 1021
Inflammation in lysosomes, elastinolytic activity in granules of human polymorphonuclear leukocytes, 1137
Inflammatory, anti-, action of 6-mercaptopurine, 785
Intestinal microvillous membranes, antibodies to, characterization and morphologic localization, 357
— microvillous membranes, antibodies to, inhibition of intrinsic factor–mediated attachment of vitamin B_{12} 375
Irradiated mice given bone marrow and thymus or thoracic duct lymphocytes, source of hemolysin-forming cells in, 821
— mice, thymus and recovery of sheep erythrocyte response in, 1327
— syngeneic mice, detection of primed cells and their proliferation upon transfer to, 171
Isoantigens of *Histocompatibility-2* and *Thymus leukemia antigen* loci, 85
Isozymes of acid phosphatase in normal and Calmette-Guérin bacillus-induced alveolar macrophages, 1031

KINETICS of renin reaction with synthetic peptide substrates, 13
— of response in mixed lymphocyte interaction, 1157
— and origin of mononuclear phagocytes, 415

LESION, initial, in allergic neuritis, phase and electron microscopic study, 469
Leukocyte antigens and mixed leukocyte culture reactivity, phenotypic expressions of major histocompatibility locus in man, 623
Leukocytes, human, polymorphonuclear, elastinolytic activity in granules of, 1137
—, polymorphonuclear, generation of factor chemotactic for, 259
—, receptors for complement on, 991
Lipopolysaccharide, endotoxic, interactions with complement system, consumption of terminal complement components, 1049
—, endotoxic, interactions with complement system, generation of chemotactic factor, 259
Loci, *Histocompatibility-2* and *Thymus leukemia*, isoantigens of, 85
Locus, major histocompatibility, in man, phenotypic expressions of, leukocyte antigens and mixed leukocyte culture reactivity, 623
—, major histocompatibility, in man, quantitative assay of antigenic disparity at, 639
Lymph nodes and spleen, distribution of antibody-forming cells of different specificities in, 729

Lymphatics, afferent, role of, in rejection of skin homografts, 197
Lymphocyte, anti-, globulin, site of action, 855
— anti-, serum, heterologous, histopathological effects of, 1281
— cultures, suppression of antiallotype-induced blast transformation in, from allotypically suppressed donors, 341
— interaction, mixed, kinetics of response, 1157
— population, small, effect of heterologous anti-lymphocyte population, 111
— responses, in vitro, abnormalities of during rubella virus infections, 47
— serum, anti-, heterologous, effect on small lymphocyte population, 111
Lymphocytes in vitro, suppression of antiallotype-induced blast transformation in lymphocyte cultures from allotypically suppressed donors, 341
—, lymphoid tissue and bone marrow, normal and immune, response to antigens in vitro, 1099
— and monocytes, interaction of in man, 1309
—, small, formation of, upon transfer of splenic white pulp to syngeneic mice, 153
—, soluble antigen and, cytotoxicity mediated by, in delayed hypersensitivity, analysis of, 1267
—, soluble antigen and, cytotoxicity mediated by, in delayed hypersensitivity, characterization of, 1237
—, soluble antigen and, cytotoxicity mediated by, in delayed hypersensitivity, correlation of in vitro response with skin reactivity, 1255
—, thoracic duct, hemolysin-forming cells in neonatally thymectomized mice reconstituted with thymus or, 801
—, thoracic duct, or bone marrow and thymus, source of hemolysin-forming cells in irradiated mice given, 821
Lymphoid cells, morphologic study of during anamnestic response, 293
— tissue and bone marrow lymphocytes, normal and immune, response to antigens in vitro, 1099
— tissue, germinal centers in, relationship to immunological memory, 153, 171
Lymphoma, Burkitt's, and infectious mononucleosis, comparison of cells and sera from, in Epstein-Barr viral and cell membrane immunofluorescence, 1021
Lymphopoiesis, thymic, and secretion by medullary epithelial cells, relation to sulfate incorporation by thymus, 927
Lysosomes, elastinolytic activity in granules of human polymorphonuclear leukocytes, 1137

MACROPHAGES, alveolar, normal and Calmette-Guérin bacillus–induced, isozymes of acid phosphatase in, 1031
—, role of in immunological maturation, 459
—, subcellular distribution of antigen in, 1059
Man, chemical characterization of glycopeptides from human γM-globulins, 699
—, digestive system of, physicochemical studies of carcinoembryonic antigens of, 387
—, elastinolytic activity in granules of polymorphonuclear leukocytes, 1137
—, interaction of monocytes and lymphocytes in, 1309
—, persistence of streptococcal Group A antibody in patients with rheumatic valvular disease, 1081
—, phenotypic expression of major histocompatibility locus in, leu-

kocyte antigens and mixed leukocyte culture reactivity, 623
Man, quantitative assay of antigenic disparity at major histocompatibility locus, 639
—, second component of human complement, its isolation, fragmentation by C'1 esterase, and incorporation into C'3 convertase, 533
—, subclass of γA-globulins (γA2) in, which lacks disulfide bonds linking heavy and light chains, 1223
—, synthesis of first component of human complement in vitro, 595
—, ultrastructure of human thyroglobulin, 1129
Marker analysis, chromosomal, of single antibody-forming cells, 839
Markers, genetic, Gm(g) and Gm(n), γG–heavy chain gene complexes with reference to, 763
Medullary epithelial cells, secretion by, and thymic lymphopoiesis, relation to sulfate incorporation by thymus, 927
Membranes, intestinal microvillous, antibodies to, characterization and morphologic localization, 357
—, intestinal microvillous, antibodies to, inhibition of intrinsic factor–mediated attachment of vitamin B_{12}, 375
Memory cells, paralysis of, X-Y-Z scheme of immunocyte maturation, 715
—, immunological, cytology of in morphologic study of lymphoid cells during anamnestic response, 293
—, immunological, relationship of germinal centers in lymphoid tissue to, 153, 171
6-Mercaptopurine, anti-inflammatory action of, 785
Metabolic responses to neonatal infection with filterable weight-depressing agent, lasting biological effects of early environmental influences, 753
Microscopy, electron, use of hybrid antibody with anti-γG and antiferritin specificities in locating cell surface antigens by, 1461
—, phase and electron, study of initial lesion in allergic neuritis, 469
Microvillous membranes, intestinal, antibodies to, characterization and morphologic localization, 357
— membranes, intestinal, antibodies to, inhibition of intrinsic factor–mediated attachment of vitamin B_{12}, 375
Monocytes and lymphocytes, interaction of in man, 1309
Mononuclear cells, chemotaxis of, 1201
— phagocytes, origin and kinetics of, 415
Mononucleosis, infectious, and Burkitt's lymphoma, comparison of cells and sera from patients with, in Epstein-Barr viral and cell membrane immunofluorescence, 1021
Morphology of lymphoid cells during anamnestic response, 293
Mucopeptide, streptococcal and staphylococcal, cross-reactivity between, 325
Mucopolysaccharide synthesis in cell cultures of chondrocytes, 1353

NEONATAL infection with filterable weight-depressing agent, metabolic responses to, lasting biological effects of, 753
— skin grafts, tolerance of H-Y antigen in mice with, 69
Neuritis, allergic, intitial lesion in, phase and electron microscopic study, 469
Neutrophils, isolation and characteri-

zation of permeability factors from, 605
Nodes, lymph, and spleen, distribution of antibody-forming cells of different specificities in, 729

OPSONIC site, primary, localization of, to Fc fragment, phagocytosis in subacute bacterial endocarditis, 553

PATHOGENETIC mechanisms in immune fever, 309
Peptide, muco-, streptococcal and staphylococcal, cross-reactivity between, 325
— substrates, synthetic, reaction of renin with, 13
Peritoneal exudate cell migration, in vitro inhibition of by chemically defined antigens, 1451
Permeability factors from neutrophils, isolation and characterization of, 605
Phagocytes, mononuclear, origin and kinetics of, 415
Phagocytosis in subacute bacterial endocarditis, localization of primary opsonic site to Fc fragment, 553
Phase and electron microscopic study of initial lesion in allergic neuritis, 469
Phenotypic expressions of major histocompatibility locus in man, leukocyte antigens and mixed leukocyte culture reactivity, 623
Phosphamide, cyclo-, graft versus host disease in mice treated with, 277
Phosphatase, acid, isozymes of, in normal and Calmette-Guérin bacillus–induced alveolar macrophages, 1031
Plaque-forming cells, antigen-stimulated proliferation of, radioautographic studies, 352
— cells, hemolytic 19S, generation time during primary and secondary response, 895
Platelet function, effect of acetylsalicylic acid on, 877
Pneumococcal strains, C-like, biological and genetic properties of, 1385
Pneumococcus, capsulation of with soluble C-like polysaccharide, biological and genetic properties of C-like pneumococcal strains, 1385
Polymorphonuclear leukocytes, human, elastinolytic activity in granules of, 1137
— leukocytes, generation of factor, chemotactic for, 259
Polysaccharide, lipo-, endotoxic, interactions with complement system, consumption of terminal complement components, 1049
—, muco-, synthesis in chondrocyte cell cultures, cellular differentiation and aging process in cartilaginous tissues, 1353
—, soluble C-like, capsulation of pneumococcus by, 1385
Precipitin and bactericidal cross-reactions among Group A streptococci, 959
Protein fragment, Bence Jones, catabolic origin of, 517
Pulp, splenic white, formation of small lymphocytes upon transfer of to syngeneic mice, 153
Purine, 6-mercapto-, anti-inflammatory action of, 785

RADIOAUTOGRAPHIC studies of plaque-forming cells, antigen-stimulated proliferation, 235
Receptors, antigen, competition of 19S and 7S, in regulation of primary immune response, 133
— for complement on leukocytes, 991

Renin, reaction with synthetic peptide substrates, 13
Rheumatic valvular disease, persistence of streptococcal Group A antibody in patients with, 1081
Rubella virus infections, abnormalities of in vitro lymphocyte responses during, 47

SECRETION by medullary epithelial cells and thymic lymphopoiesis, relation to sulfate incorporation by thymus, 927
Sensitive, antigen-, units, separate splenic, for different types of anti-sheep antibody-forming cells, 437
Sera, comparison of cells and, from patients with Burkitt's lymphoma and infectious mononucleosis, in Epstein-Barr viral and cell membrane immunofluorescence, 1021
Serum, anti-lymphocyte, heterologous, effect on small lymphocyte population, 111
—, anti-thymocyte, effects on primary and secondary responsiveness in vaccinia infection of mice, 121
—, heterologous anti-lymphocyte, histopathological effects of, 1281
— sickness–cholesterol-induced atherosclerosis, effect of removal of cholesterol diet upon, 497
Skin grafts, neonatal, tolerance of H-Y antigen in mice with, 69
— homografts, role of afferent lymphatics in rejection of, 197
— reactivity, correlation of in vitro response in cytotoxicity mediated by soluble antigen and lymphocytes in delayed hypersensitivity, 1255
Specificities, anti-γG and anti-ferritin, use of hybrid antibody with in locating cell surface antigens by electron microscopy, 1461
Specificities, different, antibody-forming cells of, distribution of in lymph nodes and spleen, 729
Specificity, competence of single cells to produce antibodies of two specificities, 223
—, individual antigenic, of antibodies to streptococcal carbohydrates, 969
— of cellular hypersensitivity, in vitro inhibition of peritoneal exudate cell migration by chemically defined antigens, 1451
Spleen cell populations separated by differential flotation in albumin gradients, analysis and in vitro response, 681
— cells, transfer of antibody response by, and linkage to major histocompatibility locus, 1
— and lymph nodes, distribution of antibody-forming cells of different specificities in, 729
Splenic antigen-sensitive units, separate, for different types of anti-sheep antibody-forming cells, 437
— white pulp, formation of lymphocytes upon transfer of, to syngeneic mice, 153
Sporobolomyces yeast, cross-reactions of an acetylphosphogalactan of, 189
Staphylococcal and streptococcal mucopeptide, immunochemical studies on cross-reactivity between, 325
Streptococcal carbohydrates, individual antigenic specificity of antibodies to, 969
— Group A antibody, persistence of, in patients with rheumatic valvular disease, 1081
— and staphylococcal mucopeptide, immunochemical studies on cross-reactivity between, 325
Streptococci, Group A, precipitin and

bactericidal cross-reactions among, 959
Streptolysin O, purified, analysis of toxic actions of, on isolated heart and separate cardiac tissues, 1401
Subcellular distribution of antigen in macrophages, 1059
Substrates, synthetic peptide, reaction of renin with, 13
Sulfate incorporation by thymus, relation to secretion by medullary epithelial cells and to thymic lymphopoiesis, 927
Syngeneic mice, detection of primed cells and their proliferation upon transfer to, 171
— mice, formation of lymphocytes upon transfer of splenic white pulp to, 153
— systems, passive transfer of tuberculin sensitivity in, 1425
Synthesis, mucopolysaccharide, in cell cultures of chondrocytes, 1353
— of first component of human complement in vitro, 595
Synthetic peptide substrates, reaction of renin with, 13

THORACIC duct lymphocytes, hemolysin-forming cells in neonatally thymectomized mice reconstituted with thymus or, 801
— duct lymphocytes or bone marrow and thymus, source of hemolysin-forming cells in irradiated mice given, 821
Thymic lymphopoiesis, and secretion by medullary epithelial cells, relation to sulfate incorporation by thymus, 927
Thymocytes, interactions of isoantigens of *Histocompatibility-2* and *Thymus leukemia antigen* loci affecting their representation on, 85
Thymus and bone marrow or thoracic duct lymphocytes, source of hemolysin-forming cells in irradiated mice given, 821
—, hemolysin-forming cells in neonatally thymectomized mice reconstituted with, or thoracic duct lymphocytes, 801
Thymus and recovery from cyclophosphamide-induced tolerance to erythrocytes, 35
— and recovery of sheep erythrocyte response in irradiated mice, 1327
—, sulfate incorporation by, relation to secretion by medullary epithelial cells and to thymic lymphopoiesis, 927
Thymus leukemia antigen (*T1a*) and *Histocompatibility-2* (*H-2*) loci, isoantigens of, 85
Thyroglobulin, human, ultrastructure of, 1129
Tolerance, cyclophosphamide-induced, to erythrocytes, thymus and recovery from, 35
— of the H-Y antigen in mice with neonatal skin grafts, 69
Toxic action of purified streptolysin O on isolated heart and separate cardiac tissues, 1401
Transfer, passive, of tuberculin sensitivity in syngeneic systems, 1425
Transplantation of mouse mammary tumor, endotoxin as adjuvator to, 1363
Tuberculin reactions, passively transferred, identification of allogeneic or hybrid cells in, by immunofluorescence, 1437
— sensitivity in syngeneic systems, passive transfer of, 1425
Tumor cells, Burkitt, comparison with infectious mononuclear cells in Epstein-Barr viral and cell membrane immunofluorescence, 1021
— cells, Burkitt, dependence of cell membrane immunofluorescence of on presence of Epstein-Barr virus, 1011
—, mouse mammary, endotoxin as adjuvator to transplantation of, 1363

VACCINIA infection, cellular immunity in, 121
Valvular disease, rheumatic, persistence of streptococcal Group A antibody in patients with, 1081
Virulence and cell wall composition in *Escherichia coli*, 399
Virus, Epstein-Barr, and cell membrane immunofluorescence, comparison of cells and sera from patients with Burkitt's lymphoma and infectious mononucleosis, 1021
—, Epstein-Barr, dependence of cell membrane immunofluorescence of Burkitt tumor cells on presence of, 1011
Virus, rubella, infections, abnormalities of in vitro lymphocyte responses during, 47
Vitamin B_{12}, inhibition of intrinsic factor–mediated attachment of in hamster brush borders, 375

WEIGHT-DEPRESSING agent, metabolic responses to neonatal infection with, lasting biological effects of early environmental influences, 753

X-Y-Z scheme of immunocyte maturation, paralysis of memory cells, 715